

# To overcome

**Current Therapeutic Problems of**

**Skin, Soft Tissue,  
Bone and Joint,  
Lymphatic Tissue,  
Respiratory and  
Surgical Infections**

**PREFER THE OPTIMAL THERAPY WITH**

**PERILACT<sup>®</sup>**  
**Cloxacillin Sodium**

# Brief Prescribing Information

## INDICATIONS

The principal indication of Cloxacillin Sodium is for the treatment of infections due to Penicillinase producing Staphylococci and other susceptible bacteria. It is used for the patients with Skin, Soft Tissue, Lymphatic Tissue, Bone, Joint and Surgical Infection.

*Bacteriologic studies to determine the causative organisms and their susceptibility to Cloxacillin Sodium is advisable.*

## CONTRAINDICATIONS

A history of a previous hypersensitivity reaction to any of the penicillins is contraindicated.

### **Pregnancy**

The safety of Cloxacillin Sodium during pregnancy has not been established.

## ADVERSE REACTIONS

Gastrointestinal disturbances such as Nausea, Epigastric Discomfort, Flatulence and Loose Stools have been noted by some patients. Skin Rashes and Allergic symptoms including Wheezing and Sneezing have occasionally been encountered. Eosinophilia with or without overt allergic manifestations has been noted in some patients during therapy.

## DOSAGE AND ADMINISTRATION

*For Bone, Respiratory and Localized Skin and Soft Tissue Infections and disseminated infections due to susceptible organisms. Adults and children weighing 20kg or more: 500mg every 6 hours before meal.*

*Children weighing less than 20kg: 50-100 mg/kg/day equally divided doses every 6 hours before meal, depending upon the severity of infections.*

## SUPPLY

**Perilact 250** capsules containing 250mg Cloxacillin as Cloxaclin Sodium in strip of 10 x 6 capsules.

**Perilact 500** capsules containing 500mg Cloxacillin as Cloxaclin Sodium in strip of 10 x 6 capsules.

# PERILACT<sup>®</sup>

## Cloxacillin Sodium


*Ideally suitable for  
infections due to  
**Penicillinase**  
producing **Staphylococci**  
and other susceptible  
microorganisms  
in different body systems.*



### पेटील्याक्ट

#### INDICATED FOR

- Osteomyelitis
- Wound infections
- Infected burns
- Furunculosis
- Boils and carbuncles
- Abscesses and cellulitis
- Otitis Media
- Pneumonia
- Tonsillitis and pharyngitis
- Staphylococcal enteritis
- Staphylococcal meningitis
- Staphylococcal Urinary Tract infections



**Prevents Penicillinase  
hydrolysis of Beta Lactam ring**

**PERILACT<sup>®</sup>**

**The brand of Cloxacillin**

**In the late 90's Staphylococcus aureus has emerged as a leading organism in wound infection with prevalence of 32% compared to 16% in 1990.**

Ref: Shrestha PC Dr., Shrestha J., Sharma VK Dr., Jan 2001, Post-Graduate Medical Journal of Nepal, Vol 2, No. 1, PP 52-55.

*For further information, please contact*



**Deurali-Janta  
Pharmaceuticals Pvt. Ltd.**

Corporate & Sales Office:  
Hattisar, Kamalpokhari,  
GPO Box 4239, Kathmandu, Nepal.  
Tel: 435167/68/69  
Fax: +977-1-435166 , 412292  
Email: djpl@ccsl.com.np



*In Technical Collaboration with:*

**RENATA LIMITED**

Bangladesh  
® Registered Trade Mark

**Deurali-Janta : Quality At Its Best**