

FORTIPLEX-m CAPSULES/DROPS

Category: Vitamins

Presentation: **Each capsule contains**

Vitamin A	BP	5000 IU
Vitamin C	BP	75mg
Vitamin D3	BP	500 IU
Vitamin E	BP	15 IU
Vitamin B1	USP	4.5mg
Vitamin B2	BP	5mg
Vitamin B6	BP	2mg
Nicotinamide	BP	45mg
Cyanocobalamin	BP	6mg
Calcium Pantothenate	BP	11mg
Folic acid	BP	1mg

Presentation: Available as 30 Capsules X 10 Blisters

INDICATION

FORTIPLEX-m is recommended for the treatment and or prevention of multivitamin deficiency. It is used as vitamin supplement during pregnancy, lactation, old-age, illness and convalescence.

DOSAGE

One capsule daily after meal or as directed by the physician

ADVERSE EFFECTS/SIDE EFFECTS

Vitamin A

- Abdominal pain.
- Appetite loss
- Bone or joint pain

Vitamin E

- Abdominal pain Discontinue
- Breast enlargement Discontinue.
- Diarrhea Discontinue.

INTERACTION

- Antacids decrease absorption of Vitamin A and fat-soluble vitamins D, E, K.
- Vitamin E: Normal amount facilitates absorption, storage in liver and utilization of Vitamin A. Excessive dosage may deplete Vitamin A stores in liver.
- Vitamin E decreases effect of iron supplement in people with iron-deficiency anemia. Iron supplements decrease Vitamin-E effect in healthy people.

- Alcohol reduces intestinal absorption of Vitamin B-1, which is necessary to metabolize alcohol.
- Aminosalicylates reduce absorption of Vitamin B-12. Antibiotics may cause false-low test results for Vitamin B-12.
- Antidepressants (tricyclic) decrease B-2 effect. Phenothiazines decrease B-2 effect.
- Estrogen or oral contraceptives increase requirements of pyridoxine. Also causes depression
- Isoniazid decreases niacin effect.

For further information, please contact:

Market Planning Department



Deurali-Janta Pharmaceuticals Pvt. Ltd.

GPO Box 4239, 355 Hattisar Road, Kamalpokhari, Kathmandu, Nepal.

Tel: 4435167/68/69 E-mail: mplanning@deuralijanta.com Website: www.deuralijanta.com